

Conversions of \mathbf{D} and \mathbf{H} between SI and Gaussian Units

Kirk T. McDonald

Joseph Henry Laboratories, Princeton University, Princeton, NJ 08544

(April 12, 2014; updated April 23, 2014)

1 Problem

The electric displacement field \mathbf{D} has dimensions charge/area and the magnetic field strength \mathbf{H} has dimensions current/length. In the SI system the units of \mathbf{D} are C(oulomb)/m² and the units of \mathbf{H} are A/m, while in the Gaussian (cgs) system the units of \mathbf{D} are sC(statcoulomb)/cm² and the units of \mathbf{H} are Oe (oersteds) where^{1,2,3}

$$1 \text{ Oe} = c_{\text{cgs}} \text{ sA(statampere)/cm} (= 1 \text{ abampere/cm in the EMU system}), \quad (1)$$

and c is the speed of light in vacuum. The conversions of coulombs to statcoulombs, of amperes to statamperes, are that [1, 10, 11]

$$1 \text{ C} = \frac{c_{\text{cgs}}}{10} \text{ sC} \approx 3 \times 10^9 \text{ sC}, \quad \text{and} \quad 1 \text{ A} = \frac{c_{\text{cgs}}}{10} \text{ sA} \approx 3 \times 10^9 \text{ sA}. \quad (2)$$

The conversion of \mathbf{D} is that [10, 11]

$$D_{\text{SI}} = 1 \text{ C/m}^2 = D_{\text{cgs}} = 4\pi c_{\text{cgs}} \times 10^{-5} \text{ sC/cm}^2 \approx 12\pi \times 10^5 \text{ sC/cm}^2, \quad (3)$$

and the conversion of \mathbf{H} is that [1, 10, 11]

$$H_{\text{SI}} = 1 \text{ A/m} = H_{\text{cgs}} = 4\pi \times 10^{-3} \text{ Oe} = 4\pi c_{\text{cgs}} \times 10^{-3} \text{ sA/cm} \approx 12\pi \times 10^7 \text{ sA/cm}. \quad (4)$$

However, from eq. (2) we find the conversions

$$1 \text{ C/m}^2 = c_{\text{cgs}} \times 10^{-5} \text{ sC/cm}^2, \quad \text{and} \quad 1 \text{ A/m} = c_{\text{cgs}} \times 10^{-3} \text{ sA/cm}. \quad (5)$$

Could it be that the conversions (3)-(4) erroneously include factors of 4π ?

This problem was suggested by Neal Carron.

¹Strictly, the dimensions of H_{cgs} follow from Ampère's law [1], $\nabla \times \mathbf{H}_{\text{cgs}} = 4\pi I_{\text{cgs}}/c_{\text{cgs}}$, as current/(length · velocity), *i.e.*, sA·s/cm². Since 1 oersted is not 1 sA·s/cm², but c_{cgs} times this, it is convenient to consider that the dimensions, cm/s, of c_{cgs} cancel a factor of s/cm in the dimensions of the oersted, leading to eq. (1). The complication due to the factors of c_{cgs} could be avoided by consideration of the EMU rather Gaussian system of units, but the issue of “extra” factors of 4π remain for this case (and for EMU units) as well.

²The oersted is sometimes defined in the EMU system as $1/4\pi$ abamp-turns/cm, meaning that a long solenoid of azimuthal current density $1/4\pi$ abamp-turns/cm produces $H_{\text{EMU}} = 1$ Oe according to Ampère's law, $\nabla \times \mathbf{H}_{\text{EMU}} = 4\pi \mathbf{J}$ in these units [1]. See, for example, [2], where the preceding is expressed somewhat indirectly. Equivalently, a long wire with current $I = 0.5$ abA produces $H_{\text{EMU}} = 2I/r = 1$ Oe = 1 abA/cm at radius $r = 1$ cm, and a loop of radius $r = 1$ cm with current $I = 1/2\pi$ abA produces $H_{\text{EMU}} = 2\pi I/r = 1$ Oe = 1 abA/cm at its center. *One should not misread abamp-turns/cm as abamp/cm to infer that 1 Oe = (1/4π) abamp/cm.*

³For a classic discussion of dimensions and units, see [3]. For amusing commentary on the oersted, see [4]. Early discussions of dimensions and units in electrodynamics include chap. X of Maxwell's Treatise [5], as well as contributions by Clausius [6], Helmholtz [7], Hertz [8] and Rücker [9].

2 Solution

2.1 Conversions of \mathbf{E} and \mathbf{B}

We first note that the conversions of the electric field \mathbf{E} and the magnetic field (or induction or flux density) \mathbf{B} do not exhibit any “extra” factors of 4π as found above for \mathbf{D} and \mathbf{H} .

The conversion of \mathbf{E} is that [1, 10, 11]

$$E_{\text{SI}} = 1 \text{ V/m} = E_{\text{cgs}} = \frac{10^6}{c_{\text{cgs}}} \text{ sV/cm}, \quad (6)$$

which is consistent with the conversion from volts to statvolts (sV) [1, 11],

$$1 \text{ V} = \frac{10^8}{c_{\text{cgs}}} \text{ sV} \approx \frac{1}{300} \text{ sV}, \quad (7)$$

The conversion of \mathbf{B} is that [1, 10, 11]

$$B_{\text{SI}} = 1 \text{ T(tesla)} = B_{\text{cgs}} = 10^4 \text{ G} = \frac{10^4}{c_{\text{cgs}}} \text{ sA/cm}, \quad (8)$$

where $1 \text{ G(gauss)} = c_{\text{cgs}} \text{ sA/cm} = 1 \text{ Oe}$. To check the consistency of eq. (8) we note that a tesla is not a primary SI unit. Using, for example, the Lorentz force law, $\mathbf{F} = q\mathbf{v} \times \mathbf{B}_{\text{SI}}$, we see that a magnetic field B_{SI} has dimensions of force/(charge · velocity) = force/(current · distance), *i.e.*, $\text{kg/A}\cdot\text{s}^2$. In Gaussian units, the force law includes a factor of $1/c_{\text{cgs}}$, so the dimensions of B_{cgs} can be considered as $(1/c_{\text{cgs}}) \text{ g/sA}\cdot\text{s}^2$ as well as a gauss, which is also equivalent to $(1/c_{\text{cgs}}) \text{ sA/cm}$. Then, the conversions

$$1 \text{ T} = 1 \frac{\text{kg}}{\text{A}\cdot\text{s}^2} = \frac{10^3 \text{ g}}{c_{\text{cgs}}/10 \text{ sA}\cdot\text{s}^2} = \frac{10^4}{c_{\text{cgs}}} \frac{\text{g}}{\text{sA}\cdot\text{s}^2} = \frac{10^4}{c_{\text{cgs}}} \text{ sA/cm} = 10^4 \text{ G}, \quad (9)$$

give a confirmation of eq. (8).

2.2 Conversions of \mathbf{D} and \mathbf{H}

In the remainder of this note we consider media of unit relative permittivity and permeability, such that $\mathbf{D}_{\text{SI}} = \epsilon_0 \mathbf{E}_{\text{SI}}$ and $\mathbf{H}_{\text{S}} = \mathbf{H}_{\text{SI}}/\mu_0$, where $\mu_0 = 4\pi \times 10^{-7} \text{ kg}\cdot\text{m/A}^2\cdot\text{s}^2 = 1/\epsilon_0 c_{\text{SI}}^2$, while $\mathbf{D}_{\text{cgs}} = \mathbf{E}_{\text{cgs}}$ and $\mathbf{B}_{\text{cgs}} = \mathbf{H}_{\text{cgs}}$.

The conversions of \mathbf{D} and \mathbf{H} differ from those for \mathbf{E} and \mathbf{B} by the factors of $\epsilon_0 = 1/c_{\text{SI}}^2 \mu_0 = 10^4/c_{\text{cgs}}^2 \mu_0 = 10^{11}/4\pi c_{\text{cgs}}^2$ and $1/\mu_0 = 10^7/4\pi$. It is these factors of 4π that make the conversions of \mathbf{D} and \mathbf{H} differ from the conversions of their units.

For example,

$$H_{\text{cgs}} = 1 \text{ Oe} = B_{\text{cgs}} = 1 \text{ G} = B_{\text{SI}} = 10^{-4} \text{ T}, \quad (10)$$

and hence,

$$H_{\text{SI}} = \frac{B_{\text{SI}}}{\mu_0} = \frac{10^3}{4\pi} \text{ A/m} = H_{\text{cgs}} = 1 \text{ Oe}, \quad (11)$$

which confirms the conversion (4).

Similarly,⁴

$$D_{\text{cgs}} = 1 \text{ sC/cm}^2 = E_{\text{cgs}} = 1 \text{ sV/cm} = E_{\text{SI}} = c_{\text{cgs}} \times 10^{-6} \text{ V/m}, \quad (12)$$

and hence,

$$D_{\text{SI}} = \epsilon_0 E_{\text{SI}} = \frac{10^{11}}{4\pi c_{\text{cgs}}^2} c_{\text{cgs}} \times 10^{-6} = \frac{10^5}{4\pi c_{\text{cgs}}} \text{ C/m}^2 = D_{\text{cgs}} = 1 \text{ sC/cm}^2, \quad (13)$$

which confirms the conversion (3).

2.3 Comments

It remains unusual that the conversions (3)-(4) include a factor of 4π times the factors associated with the conversions of the units. This occurs because \mathbf{D} and \mathbf{E} , and also \mathbf{B} and \mathbf{H} , have the same dimensions in Gaussian units⁵ but different dimensions in SI units (in that electrical current is a separate unit in the SI system but a secondary unit in the Gaussian system^{6,7,8}). The relations $\mathbf{D} = \epsilon_0 \mathbf{E} = \mathbf{E}/\mu_0 c_{\text{SI}}^2$ and $\mathbf{H} = \mathbf{B}/\mu_0$ in the SI system (in vacuum) involve factors of 4π not present in the corresponding relations ($\mathbf{D} = \mathbf{E}$ and $\mathbf{H} = \mathbf{B}$) in Gaussian units, so that while the conversions (6) and (8) of \mathbf{E} and \mathbf{B} match the conversions of their units, this cannot also hold for the conversions (3)-(4) of \mathbf{D} and \mathbf{H} .^{9,10}

⁴The dimensions in Gaussian units of the electric potential are $\text{sV} = \text{sC/cm} = \text{g}^{1/2} \text{ cm}^{1/2}/\text{s}$.

⁵As remarked in footnote 1, the dimensions of \mathbf{H} (and of \mathbf{B}) in Gaussian units are $\text{sA}\cdot\text{s/cm}^2$. Since the current in sA is charge/second, sC/s , the dimensions of \mathbf{B} and \mathbf{H} are also $\text{sC/cm}^2 = \text{sV/cm}$, the dimensions of \mathbf{D} and \mathbf{E} (in Gaussian units). And, these dimensions are also those of the gauss and the oersted. Of course, in terms of the primary units of the Gaussian system, all these equivalent dimensions are equal to $\text{g}^{1/2}/\text{cm}^{1/2}\cdot\text{s}$.

⁶“Extra” factors of 4π also occur in the conversions of \mathbf{D} and \mathbf{H} between the SI system and the so-called ESU and EMU systems of units, but not in conversions among the ESU, EMU and Gaussian systems.

⁷This issue is addressed, for example, on p. 431 of [12] where the different forms of Maxwell’s equations for \mathbf{D} , \mathbf{B} and \mathbf{H} in Gaussian and SI units (p. 427) are characterized by saying that these fields are “different physical quantities” in the two systems of units, with the implication that the conversion factors are not simply related to the conversions of the various units. Apparently, this view was adopted at an international convention on electrical units in 1930, as reported in [13, 14]. For a dissenting view, see [15].

⁸Conversions between quantities with different dimensions in different systems of units are considered somewhat abstractly in [16]. See also the much earlier paper [17].

⁹The SI system was proposed by Giorgi [18] as a “rationalized” system of units (first advocated by Heaviside [19]) in which factors of 4π would not appear explicitly in Maxwell’s equations. See also [2, 14, 20, 21]. However, factors of 4π remain present in these equations for \mathbf{B} and \mathbf{E} (contrary to the vision of Heaviside [19]) via the definitions that $\mu_0 = 4\pi \times 10^{-7}$ and $\epsilon_0 = 1/\mu_0 c_{\text{SI}}^2$. The resulting peculiar relation between \mathbf{H} (and \mathbf{D}) in “rationalized” (SI) and unrationalized (cgs) units has been discussed in [22, 23, 24].

¹⁰A fourth cgs system (in addition to Gaussian, EMU and ESU) is defined by eqs. (1-153), $\mathbf{B}_4 = \mu_0 \mathbf{H}_4$; (1-172), $\nabla \times \mathbf{B}_4 = \mu_0 \mathbf{J}$; and (1-178), $\nabla \times \mathbf{H}_4 = \mathbf{J}$, of Part 4 (pp. 4–11-14) of [25] in which the statement after eq. (1-153) that $4\pi\mu_0 = 1$ implies that the magnetic fields \mathbf{B}_4 and \mathbf{H}_4 are related by $\mathbf{B}_4 = \mathbf{B}_{\text{EMU}}/(4\pi)^2$ while $\mathbf{H}_4 = \mathbf{H}_{\text{EMU}}/4\pi = \mathbf{B}_{\text{EMU}}/4\pi = 4\pi\mathbf{B}_4$, which leads to the unusual convention (p. A-12 of [25]) that the oersted is given by $1 \text{ Oe}_4 = (1/4\pi) \text{ abA/cm}$. However, the text on p. A-12 of [25] claims that $\mu_0 = 4\pi$, in which case the equations listed above lead to a fifth version of cgs units where $\mathbf{B}_5 = \mathbf{B}_{\text{EMU}} = (4\pi)^2 \mathbf{B}_4$, while $\mathbf{H}_5 = \mathbf{H}_4 = \mathbf{H}_{\text{EMU}}/4\pi = \mathbf{B}_{\text{EMU}}/4\pi = \mathbf{B}_5/4\pi$.

References

- [1] J.D. Jackson, *Classical Electrodynamics*, 3rd ed. (Wiley, New York, 1999).
- [2] J.J. Smith, *Recommendations of IEC Technical Committee No. 24 on Electric and Magnetic Magnitudes and Units*, Trans. Am. Inst. Elec. Eng. **74**, 37 (1955),
http://physics.princeton.edu/~mcdonald/examples/EM/smith_taiee_74_37_55.pdf
- [3] R.T. Birge, *On Electric and Magnetic Units and Dimensions*, Am. Phys. Teach. **2**, 41 (1934), http://physics.princeton.edu/~mcdonald/examples/EM/birge_ajp_2_41_34.pdf
On the Establishment of Fundamental and Derived Units, with Special Reference to Electric Units, Am. Phys. Teach. **3**, 102, 171 (1935),
http://physics.princeton.edu/~mcdonald/examples/EM/birge_ajp_3_102_35.pdf
- [4] E.F. Brown, Jr, *Tutorial Paper on Dimensions and Units*, IEEE Trans. Mag. **20**, 112 (1984), http://physics.princeton.edu/~mcdonald/examples/EM/brown_ieetm_20_112_84.pdf
- [5] J.C. Maxwell, *A Treatise on Electricity and Magnetism*, vol. 2, 3rd ed. (Clarendon Press, 1892), http://physics.princeton.edu/~mcdonald/examples/EM/maxwell_treatise_v2_92.pdf
- [6] R. Clausius, *Ueber die verschiedenen Maasssysteme zur Messung electricischer und magnetischer Grössen*, Ann. d. Phys. **16**, 529 (1882),
http://physics.princeton.edu/~mcdonald/examples/EM/clausius_ap_16_529_82.pdf
Ueber den Zusammenhang zwischen den Einheiten des Magnetismus und der Electricität, Ann. d. Phys. **17**, 713 (1882),
http://physics.princeton.edu/~mcdonald/examples/EM/clausius_ap_17_713_82.pdf
The lengthy debate in the Philosophical Magazine on the above papers can be traced in http://physics.princeton.edu/~mcdonald/examples/EM/clausius_pm_13_381_82.pdf
- [7] H. Helmholtz, *Ueber absolute Maasssysteme für electriche und magnetische Grössen*, Ann. d. Phys. **17**, 42 (1882),
http://physics.princeton.edu/~mcdonald/examples/EM/helmholtz_ap_17_42_82.pdf
- [8] H. Hertz, *Ueber die Dimensionen des magnetischen Pols in verschiedenen Maasssystemen*, Ann. d. Phys. **24**, 114 (1885),
http://physics.princeton.edu/~mcdonald/examples/EM/hertz_ap_24_114_85.pdf
- [9] A.W. Rücker, *On the Suppressed Dimensions of Physical Quantities*, Proc. Phys. Soc. London **10**, 37 (1888), http://physics.princeton.edu/~mcdonald/examples/EM/rucker_pps1_10_37_88.pdf
- [10] B.W. Petley, *A Brief History of the Electrical Units to 1964*, Metrologia **31**, 481 (1995),
http://physics.princeton.edu/~mcdonald/examples/EM/petley_metrologia_31_481_95.pdf
- [11] D.J. Griffiths, *Introduction to Electrodynamics*, 4th ed., (Addison-Wesley, 2012).
- [12] R. Becker, *Electromagnetic Fields and Interactions*, (Dover Publications, 1964),
http://physics.princeton.edu/~mcdonald/examples/EM/becker_sec_g2.pdf

- [13] E. Griffiths, *Electrical and Magnetic Units*, Nature **130**, 987 (1932),
http://physics.princeton.edu/~mcdonald/examples/EM/griffiths_nature_130_987_32.pdf
- [14] A.E. Kennelly, *Notes on the July 1932 Meeting of the Committee on Symbols, Units, and Nomenclature of the International Union of Pure and Applied Physics, to Discuss Magnetic Units*, Terr. Mag. Atmos. Elec. **37**, 447 (1932),
http://physics.princeton.edu/~mcdonald/examples/EM/kennelly_tmae_37_447_32.pdf
- [15] R.G. Chambers, *Units—B, H, D, and E*, Am. J. Phys. **67**, 498 (1999),
http://physics.princeton.edu/~mcdonald/examples/EM/chambers_ajp_67_468_99.pdf
- [16] M. Kitano, *Mathematical structure of unit systems*, J. Math. Phys. **54**, 052901 (2013),
http://physics.princeton.edu/~mcdonald/examples/EM/kitano_jmp_54_052901_13.pdf
- [17] E. Weber, *A Proposal to Abolish the Absolute Electrical Unit Systems*, Trans. Am. Inst. Elec. Eng. **51**, 728 (1932),
http://physics.princeton.edu/~mcdonald/examples/EM/weber_taiee_51_728_32.pdf
- [18] G. Giorgi, *Proposals Concerning Electrical and Physics Units*, Trans. Int. Elec. Cong. vol. 1, 136 (1904), http://physics.princeton.edu/~mcdonald/examples/EM/giorgi_tiec_v1_136_04.pdf
- [19] O. Heaviside, *The Position of 4π in Electromagnetic Units*, Nature **46**, 292 (1892),
http://physics.princeton.edu/~mcdonald/examples/EM/heaviside_nature_46_292_92.pdf
- [20] A. Sommerfeld, *Über die Dimensionen der elektromagnetischen Größen*, Ann. d. Phys. **36**, 335 (1939), http://physics.princeton.edu/~mcdonald/examples/EM/sommerfeld_ap_36_335_39.pdf
- [21] A. Sommerfeld, *Electrodynamics*, (Academic Press, 1952),
http://physics.princeton.edu/~mcdonald/examples/EM/sommerfeld_em_52_secs7-8.pdf
- [22] F.S. Silsbee, *Does Rationalization Change Units?* Elec. Eng. **76**, 296 (1957),
http://physics.princeton.edu/~mcdonald/examples/EM/silsbee_ee_76_296_57.pdf
- [23] M.K. Landolt, *Concrete and Abstract Quantities and Units after Rationalization*, Elec. Eng. **80**, 690 (1961), http://physics.princeton.edu/~mcdonald/examples/EM/landolt_ee_80_690_61.pdf
- [24] F.S. Silsbee, *Systems of Electrical Units*, J. Res. Natl. Bur. Stand. **66C**, 137 (1962),
http://physics.princeton.edu/~mcdonald/examples/EM/silsbee_jrnbs_66c_137_62.pdf
- [25] E.U. Condon and H. Odishaw, eds., *Handbook of Physics*, 2nd ed., (McGraw-Hill, 1967),
http://physics.princeton.edu/~mcdonald/examples/EM/condon_handbook_67.pdf