

Supplemental Handout # 6

Symmetry Properties of Electromagnetism

The various field and source quantities, such as $\vec{E}, \vec{D}, \vec{P}, \vec{H}, \vec{B}, \vec{M}$ e.g. $\vec{J}_e, \vec{J}_m, \vec{v}, \epsilon_o, \mu_o, \vec{L}, \vec{S}, \dots$ etc. have various symmetry properties under symmetry operations such as:

$P \equiv$ Parity (Space-Inversion, $\vec{r} \rightarrow -\vec{r}$) \rightarrow Reflection in a mirror

$T \equiv$ Time Reversal (e.g. particle motion, but run backwards in time)

$C \equiv$ Electric Charge Conjugation (Charge of Particle \rightarrow Charge of Antiparticle, e.g. $e^- \rightarrow e^+$)

$M =$ Magnetic Charge Conjugation (Magnetically Charged Particle \rightarrow Magnetically charged antiparticle, e.g. $g_N \rightarrow g_S$)

The field and source quantities mentioned above fall into various generic mathematical classes of objects, or quantities:

- 1) Scalar quantities under a given symmetry transformation, designated ϕ .
- 2) Pseudoscalar quantities under a given symmetry transformation, designated p .
- 3) Polar Vector quantities under a given symmetry transformation, designated \vec{V} .
- 4) Axial, or Pseudo-Vector quantities under a given symmetry transformation, designated \vec{A} .
- 5) N^{th} rank covariant / contravariant tensors under a given symmetry transformation,

$$T_{\mu\nu}, T^{\mu\nu}, T_{\mu}^{\nu}, T_{\sigma\mu\nu}, T_{\alpha\beta\gamma}, \dots$$

Note that, e.g.:

Electric charge q is ODD under electric charge conjugation: $Ce^- = e^+$
(i.e. q behaves as a pseudoscalar quantity p under C)

Magnetic charge g_m is ODD under magnetic charge conjugation: $Mg_m^- = g_m^+$
(i.e. g_m behaves as a pseudoscalar quantity p under M)

However note also that, e.g.:

Electric charge is EVEN under magnetic charge conjugation: $Me^- = e^-$
(i.e. q behaves as a scalar quantity ϕ under M .)

Magnetic charge is EVEN under electric charge conjugation: $Cg_m^- = g_m^-$
(i.e. g_m behaves as a scalar quantity ϕ under C .)


Note also that (if \exists no magnetic charges) the combined operations $CPT = 1$ (in any order)
(i.e. $CPT =$ identity operator). If have magnetic charges, then $CPTM = 1$ (in any order).

In order to understand the distinction between Polar Vectors and Axial (or Pseudo)-Vectors under a specific symmetry transformation, consider parity P (i.e. mirror-reflection) operation:

Polar Vector \vec{V}

e.g. direction vector \vec{r}
 P , Parity is space-inversion

i.e. $\vec{r} \rightarrow -\vec{r}$
 $x \rightarrow -x$
 $y \rightarrow -y$
 $z \rightarrow -z$




\parallel -component of \vec{V}
 unchanged under Parity
 \perp -component of \vec{V}
 changes sign under Parity

i.e. $PV_{\parallel} = V_{\parallel}$
 $PV_{\perp} = -V_{\perp}$


Axial (or Pseudo) Vector \vec{A}

e.g. spinning top –
 angular momentum \vec{L}




\parallel -components of \vec{A}
 reversed under Parity
 \perp -components of \vec{A}
 unchanged under Parity

i.e. $PA_{\parallel} = -A_{\parallel}$
 $PA_{\perp} = A_{\perp}$




Time Reversal:

Particle
 Moving with
 Velocity \vec{v}




Time-reversed
 situation:



$\vec{v}' = T\vec{v} = -\vec{v}$ (velocity is odd under Time reversal)


Electric Current Flowing in a Long Wire:



$$\begin{array}{l} \therefore \left. \begin{array}{l} T \vec{v}_d = -\vec{v}_d \\ T \vec{J}_e^{free} = -\vec{J}_e^{free} \\ T I = -I \\ T \vec{B} = -\vec{B} \\ T \vec{H} = -\vec{H} \\ T \vec{M} = -\vec{M} \end{array} \right\} \begin{array}{l} \text{all} \\ \text{odd} \\ \text{under} \\ T \end{array} \end{array} \quad \begin{array}{l} \vec{J}_e^{free} = n_e q \vec{v}_d \\ I = \vec{J}_e^{free} \cdot \vec{A} \\ \vec{B} = \frac{\mu_0}{4\pi} \int \frac{\vec{I} \times d\vec{\ell}}{r^2} \\ \vec{B} = \mu \vec{H} \\ \vec{B} = \mu_0 \vec{H} + \vec{M} \end{array} \quad \begin{array}{l} T e^- = e^- \text{ (} q \text{ is even under } T \text{)} \\ T \mu = \mu \text{ (} \mu \text{ is even under } T \text{)} \end{array}$$

Current Flowing in a Loop:


$$\begin{array}{l} \therefore T \vec{m} = -\vec{m} \quad (\text{mag. dipole moment}) \\ \text{And } T \vec{M} = -\vec{M} \quad (\text{magnetization}) \end{array}$$




By considering a parallel-plate capacitor, it can be seen that \vec{E}, \vec{D} and \vec{P} fields, \vec{p} = electric dipole moment, ϵ = permittivity, etc. are all even under time reversal.

$$\begin{array}{l} T \vec{E} = \vec{E} \quad (\text{e.g. } \vec{E} = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \hat{r}) \quad T q = +q \text{ (even under } T \text{)} \\ T \vec{P} = \vec{P} \\ T \vec{D} = \vec{D} = \epsilon \vec{E} \quad T \epsilon = \epsilon \end{array}$$


Parity and Magnetic Fields




Electric Charge Conjugation & Magnetic Fields



Time Reversal & Magnetic Fields



Magnetic Charge Conjugation & Magnetic Fields



Summary of Symmetry Properties of Kinematic & Electromagnetic Quantities

$\phi \equiv$ Scalar Quantity $p \equiv$ Pseudoscalar Quantity $\vec{V} =$ Polar Vector $\vec{A} =$ Axial Vector (Pseudo-Vector)

	KINEMATIC AND/OR ELECTROMAGNETIC QUANTITY	PARITY (SPACE INVERSION) $P(\vec{r} \rightarrow -\vec{r})$	CHARGE (ELECTRIC) CONJUGATION $C(q_e \rightarrow -q_e)$	TIME REVERSAL $T(t \rightarrow -t)$	MAGNETIC CHARGE CONJUGATION $M(q_m \rightarrow -q_m)$
KINEMATICAL	\vec{r}	-	\vec{V}	+	\vec{V}
	$\vec{v} = \frac{d\vec{r}}{dt}; \vec{p} = m\vec{v}$	-	\vec{V}	+	\vec{V}
	$\vec{a} = \frac{d\vec{v}}{dt} = \frac{d^2\vec{r}}{dt^2}; \vec{F} = m\vec{a}$	-	\vec{V}	+	\vec{V}
	$\vec{L} = \vec{r} \times \vec{p}$	+	\vec{A}	-	\vec{A}
	$\vec{\tau} = \vec{r} \times \vec{F}$	+	\vec{A}	-	\vec{A}
	KE. = $\frac{1}{2}mv^2$, PE. = $\frac{1}{2}kx^2$ WORK = $\vec{F} \cdot \vec{x}$	+	ϕ	+	ϕ
E+M CHARGES	$e, \lambda_e, \sigma_e, \rho_e$ (eg. $q_m = nh$)	+	ϕ	-	p
	$q_m, \lambda_m, \sigma_m, \rho_m$ $q_m = q_e v$	-	p	+	ϕ
VACUUM DIELECTRIC/MAGNETIC MATERIALS	$\epsilon_0, \epsilon, \chi_e, K_e = \frac{\epsilon}{\epsilon_0} = \chi_e + 1$	+	ϕ	+	ϕ
	$\mu_0, \mu, \chi_m, K_m = \frac{\mu}{\mu_0} = \chi_m + 1$	+	ϕ	+	ϕ
E+M CURRENTS	$\vec{I}_e, \vec{K}_e, \vec{J}_e = n_e q_e \vec{v}$	-	\vec{V}	-	\vec{V}
	$\vec{I}_m, \vec{K}_m, \vec{J}_m = n_m q_m \vec{v}$	-	\vec{A}	+	\vec{A}
ELECTRIC FIELD QUANTITIES	\vec{V}_E	-	ϕ	+	ϕ
	GRADIENT CURV. DIVERGENCE $\vec{\nabla} = \frac{\partial}{\partial x}\hat{x} + \frac{\partial}{\partial y}\hat{y} + \frac{\partial}{\partial z}\hat{z}$	+	\vec{V}	+	\vec{V}
	$\vec{E} = -\vec{\nabla}V_E = \frac{1}{4\pi\epsilon_0} \frac{q_e}{r^2} \hat{r}$	-	\vec{V}	-	\vec{V}
	$\vec{D} = \epsilon \vec{E}$	-	\vec{V}	+	\vec{V}
	$\vec{P} = \vec{D} - \epsilon_0 \vec{E} = \epsilon_0 \chi_e \vec{E}; \vec{p} = q_e \vec{d}$	-	\vec{V}	-	\vec{V}
MAGNETIC FIELD QUANTITIES	$\vec{B} = \frac{\mu_0}{4\pi} \frac{q_m}{r^2} \hat{r}, \frac{\mu_0}{4\pi} I_e \oint_c \frac{d\vec{x} \times \hat{r}}{r^2}, \vec{\nabla} \times \vec{A}$	-	\vec{A}	+	\vec{A}
	$\vec{H} = \frac{1}{\mu} \vec{B} = -\vec{\nabla}V_m$	-	\vec{A}	-	\vec{A}
	$\vec{M} = \vec{H} - \frac{1}{\mu_0} \vec{B} = \chi_m \vec{H}, \vec{m} = I_e \vec{a} = q_m \vec{d}$	-	\vec{A}	-	\vec{A}
	$\vec{A} = \frac{\mu_0}{4\pi} I_e \oint_c \frac{d\vec{l}}{r}$ MAGNETIC VECTOR POTENTIAL	-	\vec{A}	+	\vec{A}
	V_m	-	p	-	p
MAXWELL'S EQUATIONS	$\vec{\nabla} \cdot \vec{D} = \rho_e^{free}$	+	ϕ	-	p
	$\vec{\nabla} \cdot \vec{B} = \rho_m^{free}$	-	p	+	ϕ
	$\vec{\nabla} \times \vec{E} = -\vec{J}_m^{free} - \frac{\partial \vec{B}}{\partial t}$	-	\vec{A}	+	\vec{A}
	$\vec{\nabla} \times \vec{H} = \vec{J}_e^{free} + \frac{\partial \vec{D}}{\partial t}$	-	\vec{V}	-	\vec{V}
EM ENERGY FLOW, etc.	$\vec{S} = \vec{E} \times \vec{H}$ POYNTING'S VECTOR	+	\vec{V}	-	\vec{V}
	$T_{\mu\nu}$ MAXWELL'S STRESS TENSOR (FRANK)	+	\vec{V}	+	\vec{V}